

Black-Scholes(-Merton)*

Preliminary draft — please do not cite without permission

Gabriel E. Cabrera Guzmán[†] 

This revision: 21 August 2026

Abstract

These notes derive the Black-Scholes-Merton partial differential equation and its European-call solution from first principles. We begin with the Taylor series expansion, including a geometric interpretation of the second-order remainder, and pass through Itô's calculus, the geometric-Brownian-motion assumption, and Itô's lemma before constructing the delta-hedged portfolio that yields the Black-Scholes PDE. We conclude by evaluating the risk-neutral expectation that produces the closed-form call price $C = S_t N(d_1) - Ke^{-r(T-t)} N(d_2)$ and by showing that S_T is lognormally distributed under $\mathbb{Q}$.

Keywords: Black-Scholes, Itô's lemma, Taylor expansion, option pricing, risk-neutral valuation.

JEL Classification: G12, G13, C60.

*These are working note. Any errors are my own.

[†]Email: gabriel.cabreraguzman@postgrad.manchester.ac.uk. Address: The University of Manchester.

1. Introduction

The Black-Scholes-Merton framework ([Black and Scholes 1973](#); [Merton 1973](#)) is the canonical no-arbitrage model for pricing European options on a non-dividend-paying stock. Its two workhorses are the Taylor expansion — which controls how a smooth function responds to small perturbations in its arguments — and Itô’s calculus ([Itô 1951](#)), which supplies the additional correction terms that appear when one of those arguments is a diffusion. Combining the two, a self-financing portfolio short one option and long $\partial V / \partial S$ shares of the underlying turns out to be locally deterministic, so it must earn the risk-free rate. That local no-arbitrage condition is the Black-Scholes partial differential equation (PDE). Standard textbook treatments ([Hull 2021](#); [Shreve 2004](#)) fill in the technical background.

The remainder of this note is organised as follows. Section [2](#) develops the Taylor expansion and its geometric interpretation. Section [3](#) introduces the Wiener process and the geometric-Brownian-motion (GBM) dynamics for the underlying. Section [4](#) proves Itô’s lemma for a function of a single GBM. Section [5](#) constructs the delta-hedged portfolio and derives the Black-Scholes PDE, evaluates the risk-neutral expectation to obtain the closed-form call price, and shows that S_T is lognormally distributed under $\mathbb{Q}$. Section [6](#) concludes.

2. Taylor Series Expansion

A function $f(x + \Delta x)$ may be Taylor-expanded about a base point x whenever $f(\cdot)$ is sufficiently differentiable in a neighbourhood of x . The value at the nearby point $x + \Delta x$ is expressed in terms of the value of the function and its derivatives at x , with higher-order

derivatives providing increasingly accurate adjustments as $\Delta x \rightarrow 0$:

$$\begin{aligned} f(x + \Delta x) = f(x) &+ \frac{df(x)}{dx} \Delta x + \frac{d^2 f(x)}{dx^2} \frac{\Delta x^2}{2!} + \frac{d^3 f(x)}{dx^3} \frac{\Delta x^3}{3!} \\ &+ \dots + \frac{d^N f(x)}{dx^N} \frac{\Delta x^N}{N!} + \dots \end{aligned} \quad (1)$$

Equivalently, $f(x)$ may be expanded about an arbitrary base point a :

$$\begin{aligned} f(x) = f(a) &+ \frac{df(a)}{dx} (x - a) + \frac{d^2 f(a)}{dx^2} \frac{(x - a)^2}{2!} + \frac{d^3 f(a)}{dx^3} \frac{(x - a)^3}{3!} \\ &+ \dots + \frac{d^N f(a)}{dx^N} \frac{(x - a)^N}{N!} + \dots \end{aligned} \quad (2)$$

The two forms are the same identity; the second is convenient when a is a natural expansion point (e.g., the origin for Maclaurin series) and $(x - a)$ need not be small.

2.1 The case of e^x

Let $f(x) = e^x$, for which every derivative satisfies

$$\frac{d^k}{dx^k} e^x = e^x \quad \Rightarrow \quad \frac{d^k}{dx^k} e^0 = 1.$$

Taking $\Delta x = 0.2$ in Eq. (1) the third-order Taylor polynomial is

$$e^{x+0.2} \approx e^x + e^x(0.2) + \frac{e^x}{2!}(0.2)^2 + \frac{e^x}{3!}(0.2)^3 \equiv P_3(\Delta x).$$

This approximation uses only the function and its derivatives at x up to third order, and it becomes increasingly accurate as $\Delta x \rightarrow 0$. The progression from a zeroth-order (constant) approximation to a third-order (cubic) approximation is shown in Figure 1.

Alternatively, expanding e^x about $a = 0$ (the Maclaurin series) and truncating at third order gives

$$e^x \approx 1 + \frac{x}{1!} + \frac{x^2}{2!} + \frac{x^3}{3!} \equiv P_3(x),$$




Figure 1. Taylor expansion of $e^{x+\Delta x}$ about the base point $x = 0$.

Note: This figure plots the Taylor expansion of $e^{x+\Delta x}$ about $x = 0$ at successively higher orders. In each panel the blue solid curve is the exact function e^x , the orange dashed curve is the Taylor polynomial $P_n(\Delta x)$ truncated at order n , the red dot marks the base point $(0, 1)$, and the purple dot marks the target $(\Delta x, e^{\Delta x})$ with $\Delta x = 0.7$. The dotted grey step shows the horizontal displacement Δx and the vertical rise $e^{\Delta x} - 1$ that the polynomial has to close as n grows.

which produces a finite-order approximation of e^x near the expansion point. As n grows, $P_n(x)$ approximates e^x increasingly well:

- **0th order (constant):** $P_0(x) = 1$
- **1st order (tangent line):** $P_1(x) = 1 + x$
- **2nd order (adds curvature):** $P_2(x) = 1 + x + x^2/2!$
- **3rd order and higher:** progressively improve local accuracy.

The progression is illustrated in Figure 2.

2.2 Geometric interpretation of the second-order Taylor term

The second-order term $\frac{1}{2}f''(a)(x-a)^2$ is often dismissed as an algebraic afterthought, but it has a crisp geometric reading that makes Itô's lemma much easier to remember. The trick is to Taylor-expand not f itself but the *area* under f : doing so turns every term of the expansion into a shape you can point at.

Fix a left endpoint x_0 and define the running area

$$F(x) \equiv \int_{x_0}^x f(t) dt, \quad (3)$$

so that $F(x)$ measures the signed area under f from x_0 out to the moving right endpoint x . By the fundamental theorem of calculus,

$$F'(x) = f(x), \quad F''(x) = f'(x),$$

that is, the first derivative of the area is the *height* of the curve and the second derivative is the *slope* of the curve. Substituting into the second-order Taylor expansion of F about the base point a gives




Figure 2. Maclaurin polynomials $P_n(x)$ approximating e^x .

Note: This figure plots the Maclaurin polynomials $P_n(x)$ approximating e^x on the interval $x \in [-1, 2]$. The blue solid curve is the exact function e^x , the orange dashed curve is $P_n(x)$ truncated at order n , and the red dot marks the common expansion point $(0, 1)$. As n increases the dashed curve tracks e^x over a widening neighbourhood of the origin, illustrating that each additional term extends the range on which the polynomial approximation is accurate.

$$F(x) \approx F(a) + f(a)(x-a) + \frac{1}{2} f'(a)(x-a)^2. \quad (4)$$

Each term on the right-hand side is a shape visible in Figure 3:

1. **Zeroth order:** $F(a)$ (**blue region**). The area already accumulated up to the base point a . It is the value of the area function *at* a , before the endpoint moves.
2. **First order:** $f(a)(x-a)$ (**gold rectangle**). If we naively pretend that f is constant and equal to $f(a)$ on $[a, x]$, the extra area swept out as the endpoint moves from a to x is exactly a rectangle of height $f(a)$ and base $(x-a)$. This is the *linear* estimate of the increment $F(x) - F(a)$ — what a first-order Taylor expansion of F contributes.
3. **Second order:** $\frac{1}{2} f'(a)(x-a)^2$ (**pink triangle**). The rectangle is systematically wrong whenever f is not flat: it *underestimates* the increment when f is rising ($f'(a) > 0$) and *overestimates* it when f is falling. Locally the curve is best replaced by its tangent line at a , which lifts above the top of the rectangle by $f'(a)(x-a)$ at the right endpoint. The triangle wedged between the top of the rectangle and the tangent line has base $(x-a)$ and height $f'(a)(x-a)$, so its area is

$$\frac{1}{2} \underbrace{(x-a)}_{\text{base}} \underbrace{f'(a)(x-a)}_{\text{height}} = \frac{1}{2} f'(a)(x-a)^2,$$

exactly the second-order term in Eq. (4).

In both panels of the figure the pink triangle is bounded above by the tangent line, and across $[a, x]$ the drawn curve lies along that tangent — which is exactly the substitution a

second-order expansion makes. What it leaves out shows up just past x , where the drawn curve bends away above the tangent and the two visibly separate. That widening gap holds everything left over — the cubic, quartic, and higher-order Taylor terms. At the step drawn in the figure the three pieces are correctly ordered: the triangle is 43% of the rectangle, and the sliver is 6% of the whole increment. As the step $(x - a)$ shrinks that ordering only sharpens, because the sliver vanishes faster than the pink triangle, which in turn vanishes faster than the gold rectangle. The triangle is therefore the dominant correction to the rectangle once linear information has been exhausted.

Two takeaways deserve emphasis. First, the second-order term is *not* a rounding error but the **leading-order mismatch** between the tangent line and a curve of nonzero curvature, and its size grows quadratically with the step $(x - a)$. Second, this same accounting argument — a linear “tangent” prediction plus a quadratic correction whose sign is set by the local curvature — reappears verbatim when we Taylor-expand the derivative price $V(S_t, t)$ in Section 4. There the tangent piece is the option’s delta, $\partial V / \partial S$, and the quadratic correction is exactly the $\frac{1}{2} \sigma^2 S_t^2 \partial^2 V / \partial S^2$ gamma term that survives Itô’s rules. The geometric picture is the same; only the object being expanded changes.




Figure 3. Geometric interpretation of the second-order Taylor expansion of the area function

Note: The area function is $F(x) = \int_{x_0}^x f(t) dt$, and blue is $F(a)$, the area already accumulated at the base point. The increment $F(x) - F(a)$ splits into a gold rectangle of area $f(a)(x-a)$ and a pink triangle of area $\frac{1}{2}f'(a)(x-a)^2$ — the linear and quadratic terms of the expansion. Across $[a, x]$ the black line is drawn as the tangent at a , the substitution a second-order expansion makes, so there the increment is exactly rectangle plus triangle. Past x the line bends away from the tangent, and that gap is the higher-order remainder. The dashed box marks the interval the inset magnifies. At the step drawn, the triangle is 43% of the rectangle and the remainder 6% of the increment.

3. Itô's Process

Consider a smooth function V depending on the state S_t and on time t . Applying a second-order Taylor expansion as in Eq. (1) to $V(S_t + \Delta S_t, t + \Delta t)$ about (S_t, t) yields

$$\begin{aligned}
V(S_t + \Delta S_t, t + \Delta t) \approx & V(S_t, t) + \frac{\partial V}{\partial S} \Delta S_t + \frac{\partial V}{\partial t} \Delta t + \frac{1}{2} \frac{\partial^2 V}{\partial S^2} (\Delta S_t)^2 \\
& + \frac{\partial^2 V}{\partial S \partial t} \Delta S_t \Delta t + \frac{1}{2} \frac{\partial^2 V}{\partial t^2} (\Delta t)^2,
\end{aligned}$$

where all partial derivatives are evaluated at (S_t, t) and higher-order terms are of smaller order (they will vanish once we apply Itô's rules below). Define the increment

$$\Delta V_t := V(S_t + \Delta S_t, t + \Delta t) - V(S_t, t).$$

Then

$$\Delta V_t \approx \frac{\partial V}{\partial S} \Delta S_t + \frac{\partial V}{\partial t} \Delta t + \frac{1}{2} \frac{\partial^2 V}{\partial S^2} (\Delta S_t)^2 + \frac{\partial^2 V}{\partial S \partial t} \Delta S_t \Delta t + \frac{1}{2} \frac{\partial^2 V}{\partial t^2} (\Delta t)^2.$$

Formally replacing the finite increments by differentials,

$$dV_t \approx \frac{\partial V}{\partial S} dS_t + \frac{\partial V}{\partial t} dt + \frac{1}{2} \frac{\partial^2 V}{\partial S^2} (dS_t)^2 + \frac{\partial^2 V}{\partial S \partial t} dS_t dt + \frac{1}{2} \frac{\partial^2 V}{\partial t^2} (dt)^2. \quad (5)$$

Up to this point the derivation is an ordinary Taylor expansion. Stochastic calculus enters when we specify the dynamics of S_t and apply Itô's rules to the differentials dS_t , dW_t , and dt .

3.1 Geometric Brownian motion

Assume that the underlying S_t (e.g., a stock price) follows the diffusion

$$dS_t = \mu S_t dt + \sigma S_t dW_t, \quad (6)$$

where μ is the drift (the instantaneous expected rate of return) and σ is the volatility, so that σS_t is the instantaneous volatility of the stock price over the short interval dt . The

key stochastic ingredient is dW_t , where $(W_t)_{t \geq 0}$ is a standard Wiener process.

3.2 The Wiener process

The Wiener process $(W_t)_{t \geq 0}$ (also called standard Brownian motion) is a continuous-time Markov stochastic process with zero drift and unit variance rate. Formally, it is characterised by two defining properties.

1. *Normal increments.* Over a small time interval dt , the increment satisfies

$$dW_t = \epsilon \sqrt{dt}, \quad \epsilon \sim \mathcal{N}(0, 1).$$

2. *Independent increments.* The increments dW_t over non-overlapping intervals dt are mutually independent.

The first property implies that dW_t is normally distributed with

- $\mathbb{E}[dW_t] = \sqrt{dt} \mathbb{E}[\epsilon] = 0$,
- $\text{Var}(dW_t) = dt \text{Var}(\epsilon) = dt$,
- $\sqrt{\text{Var}(dW_t)} = \sqrt{dt}$.

The second property implies the Markov property: only the current value of the process is relevant for forecasting future increments.

4. Itô's Lemma

Substituting Eq. (6) into Eq. (5) and applying the Itô rules

$$(dt)^2 = 0, \quad (dW_t)^2 = dt, \quad dt dW_t = 0,$$

we obtain

$$\begin{aligned}
dV_t &= \frac{\partial V}{\partial S} dS_t + \frac{\partial V}{\partial t} dt + \frac{1}{2} \frac{\partial^2 V}{\partial S^2} (dS_t)^2 \\
&= \frac{\partial V}{\partial S} (\mu S_t dt + \sigma S_t dW_t) + \frac{\partial V}{\partial t} dt + \frac{1}{2} \frac{\partial^2 V}{\partial S^2} (\mu S_t dt + \sigma S_t dW_t)^2 \\
&= \frac{\partial V}{\partial S} (\mu S_t dt + \sigma S_t dW_t) + \frac{\partial V}{\partial t} dt + \frac{1}{2} \frac{\partial^2 V}{\partial S^2} (\mu^2 S_t^2 (dt)^2 + 2\mu\sigma S_t^2 dt dW_t + \sigma^2 S_t^2 (dW_t)^2) \\
&= \frac{\partial V}{\partial S} (\mu S_t dt + \sigma S_t dW_t) + \frac{\partial V}{\partial t} dt + \frac{1}{2} \sigma^2 S_t^2 \frac{\partial^2 V}{\partial S^2} dt \\
&= \left(\frac{\partial V}{\partial S} \mu S_t + \frac{\partial V}{\partial t} + \frac{1}{2} \frac{\partial^2 V}{\partial S^2} \sigma^2 S_t^2 \right) dt + \frac{\partial V}{\partial S} \sigma S_t dW_t.
\end{aligned} \tag{7}$$

This is the standard form of Itô's lemma for a function of a single stochastic variable S_t . Comparing Eq. (7) with the naive chain-rule expansion, the additional $\frac{1}{2}\sigma^2 S_t^2 \partial^2 V / \partial S^2 dt$ term is exactly the Itô correction. Geometrically, it plays the same role as the pink triangle in Section 2.2: a quadratic correction whose sign is set by the local curvature of V in S .

5. Black-Scholes(-Merton)

5.1 The delta-hedged portfolio and the PDE

The discrete-time version of Eq. (7) is

$$\Delta V_t = \left(\frac{\partial V}{\partial S} \mu S_t + \frac{\partial V}{\partial t} + \frac{1}{2} \frac{\partial^2 V}{\partial S^2} \sigma^2 S_t^2 \right) \Delta t + \frac{\partial V}{\partial S} \sigma S_t \Delta W_t, \tag{8}$$

where $\Delta W_t = \epsilon \sqrt{\Delta t}$ with $\epsilon \sim \mathcal{N}(0, 1)$. It follows that a portfolio short one derivative and long $\partial V / \partial S$ shares of the underlying can be constructed so that the Wiener term cancels. Concretely, define

$$\begin{aligned}
&-1 : \text{ derivative} \\
&+ \frac{\partial V}{\partial S} : \text{ shares of the stock,}
\end{aligned}$$

so that the value of the portfolio at time t is

$$\Pi_t = -V(S_t, t) + \frac{\partial V(S_t, t)}{\partial S} S_t.$$

Over the short interval Δt its change is

$$\Delta \Pi_t = -\Delta V_t + \frac{\partial V}{\partial S} \Delta S_t,$$

where

$$\Delta S_t = \mu S_t \Delta t + \sigma S_t \Delta W_t.$$

Substituting Eq. (8) for ΔV_t and the expression for ΔS_t ,

$$\begin{aligned} \Delta \Pi_t &= -\left(\frac{\partial V}{\partial S} \mu S_t + \frac{\partial V}{\partial t} + \frac{1}{2} \frac{\partial^2 V}{\partial S^2} \sigma^2 S_t^2\right) \Delta t - \frac{\partial V}{\partial S} \sigma S_t \Delta W_t \\ &\quad + \frac{\partial V}{\partial S} (\mu S_t \Delta t + \sigma S_t \Delta W_t) \\ &= -\left(\frac{\partial V}{\partial t} + \frac{1}{2} \frac{\partial^2 V}{\partial S^2} \sigma^2 S_t^2\right) \Delta t. \end{aligned}$$

The ΔW_t terms cancel, so over the interval Δt the portfolio is *locally riskless*: its change does not depend on ΔW_t . By no arbitrage it must earn the risk-free rate r :

$$\Delta \Pi_t = r \Pi_t \Delta t = r \left(-V + \frac{\partial V}{\partial S} S_t\right) \Delta t.$$

Equating the two expressions for $\Delta \Pi_t$ and dividing by Δt yields

$$-\left(\frac{\partial V}{\partial t} + \frac{1}{2} \frac{\partial^2 V}{\partial S^2} \sigma^2 S_t^2\right) = r \left(-V + \frac{\partial V}{\partial S} S_t\right),$$

which rearranges to the **Black-Scholes PDE**:

$$\frac{\partial V}{\partial t} + \frac{1}{2} \sigma^2 S^2 \frac{\partial^2 V}{\partial S^2} + r S \frac{\partial V}{\partial S} - r V = 0. \quad (9)$$

Eq. (9) has many solutions, one for each choice of boundary conditions. For a European call struck at K and maturing at T , the terminal condition is

$$V(S_T, T) = \max(S_T - K, 0),$$

and the PDE together with this boundary condition admits the closed-form solution

$$C(S_t, t) = S_t N(d_1) - Ke^{-r(T-t)} N(d_2), \quad (10)$$

with

$$d_1 = \frac{\ln(S_t/K) + (r + \frac{1}{2}\sigma^2)(T-t)}{\sigma\sqrt{T-t}}, \quad d_2 = d_1 - \sigma\sqrt{T-t},$$

where $N(\cdot)$ is the standard normal CDF.

5.2 Where do $N(d_1)$ and $N(d_2)$ come from?

The hedging argument shows that the value $V(S, t)$ of any derivative written on S_t must satisfy the Black-Scholes PDE Eq. (9) together with the terminal condition imposed by the payoff. A key result from stochastic analysis, the **Feynman-Kac theorem** (Shreve 2004), states that the solution can be expressed as a discounted expectation under the risk-neutral measure $\mathbb{Q}$ (Harrison and Pliska 1981). For a European call,

$$V(S_t, t) = e^{-r(T-t)} \mathbb{E}^{\mathbb{Q}}[(S_T - K)^+ | S_t],$$

where under $\mathbb{Q}$ the stock evolves according to

$$dS_t = rS_t dt + \sigma S_t dW_t^{\mathbb{Q}}.$$

Solving this SDE (see Section 5.3) shows that S_T is lognormally distributed:

$$S_T = S_t \exp\left((r - \frac{1}{2}\sigma^2)(T - t) + \sigma\sqrt{T - t}Z\right), \quad Z \sim N(0, 1).$$

Writing the payoff as $(S_T - K)^+ = (S_T - K)\mathbf{1}_{\{S_T > K\}}$, the event $\{S_T > K\}$ becomes a linear inequality in the standard normal variable Z , and the expectation reduces to two Gaussian integrals:

$$\begin{aligned} V(S_t, t) &= e^{-r(T-t)} \mathbb{E}^{\mathbb{Q}}[(S_T - K)^+ \mid S_t] \\ &= e^{-r(T-t)} \underbrace{(\mathbb{E}^{\mathbb{Q}}[S_T \mathbf{1}_{\{S_T > K\}} \mid S_t])}_A - K \underbrace{\mathbb{Q}(S_T > K \mid S_t)}_B. \end{aligned}$$

Solving for B . Using the explicit form of S_T ,

$$\begin{aligned} \mathbb{Q}(S_T > K \mid S_t) &= \mathbb{Q}\left(S_t \exp\left((r - \frac{1}{2}\sigma^2)(T - t) + \sigma\sqrt{T - t}Z\right) > K\right) \\ &= \mathbb{Q}\left(Z > \frac{\ln(K/S_t) - (r - \frac{1}{2}\sigma^2)(T - t)}{\sigma\sqrt{T - t}}\right) \\ &= \mathbb{Q}\left(Z > -\frac{\ln(S_t/K) + (r - \frac{1}{2}\sigma^2)(T - t)}{\sigma\sqrt{T - t}}\right) \\ &= \mathbb{Q}(Z > -d_2). \end{aligned}$$

Since $Z \sim N(0, 1)$ is symmetric about zero,

$$B = \mathbb{Q}(S_T > K \mid S_t) = \mathbb{Q}(Z > -d_2) = \mathbb{Q}(Z < d_2) = N(d_2).$$

Solving for A .

$$\begin{aligned} \mathbb{E}^{\mathbb{Q}}[S_T \mathbf{1}_{\{S_T > K\}} \mid S_t] &= \mathbb{E}^{\mathbb{Q}}\left[S_t \exp\left((r - \frac{1}{2}\sigma^2)(T - t) + \sigma\sqrt{T - t}Z\right) \mathbf{1}_{\{S_T > K\}} \mid S_t\right] \\ &= S_t e^{(r - \frac{1}{2}\sigma^2)(T - t)} \mathbb{E}^{\mathbb{Q}}[e^{\sigma\sqrt{T - t}Z} \mathbf{1}_{\{Z > -d_2\}}]. \end{aligned}$$

The Gaussian integral

$$\mathbb{E}[e^{aZ} \mathbf{1}_{\{Z > -d_2\}}] = \int_{-d_2}^{\infty} e^{az} \phi(z) dz, \quad a = \sigma\sqrt{T-t},$$

is evaluated by completing the square in the exponent. This yields

$$\mathbb{E}[e^{aZ} \mathbf{1}_{\{Z > -d_2\}}] = e^{\frac{1}{2}a^2} N(d_1),$$

so that

$$A = S_t e^{r(T-t)} N(d_1).$$

Substituting A and B back into the valuation formula reproduces the Black-Scholes call price:

$$C(S_t, t) = S_t N(d_1) - K e^{-r(T-t)} N(d_2),$$

which matches Eq. (10). The put price follows by the same logic (or by put-call parity):

$$P(S_t, t) = K e^{-r(T-t)} N(-d_2) - S_t N(-d_1).$$

5.3 Why is S_T lognormally distributed?

To solve the SDE under $\mathbb{Q}$,

$$dS_t = rS_t dt + \sigma S_t dW_t,$$

consider the process $X_t = \ln S_t$. Applying Itô's lemma to $f(s) = \ln s$, for which

$$f'(S_t) = \frac{1}{S_t}, \quad f''(S_t) = -\frac{1}{S_t^2},$$

we obtain

$$dX_t = f'(S_t) dS_t + \frac{1}{2} f''(S_t) (dS_t)^2.$$

Substituting $dS_t = rS_t dt + \sigma S_t dW_t$ and using the Itô rules $(dW_t)^2 = dt$, $(dt)^2 = 0$, $dt dW_t = 0$,

$$\begin{aligned} dX_t &= \frac{1}{S_t} (rS_t dt + \sigma S_t dW_t) + \frac{1}{2} \left(-\frac{1}{S_t^2} \right) \sigma^2 S_t^2 dt \\ &= r dt + \sigma dW_t - \frac{1}{2} \sigma^2 dt \\ &= \left(r - \frac{1}{2} \sigma^2 \right) dt + \sigma dW_t. \end{aligned}$$

Integrating from t to T ,

$$\int_t^T dX_s = \int_t^T \left(r - \frac{1}{2} \sigma^2 \right) ds + \int_t^T \sigma dW_s,$$

the left-hand side equals $X_T - X_t = \ln S_T - \ln S_t$, so

$$\ln S_T = \ln S_t + \left(r - \frac{1}{2} \sigma^2 \right) (T - t) + \sigma (W_T - W_t).$$

Exponentiating,

$$S_T = S_t \exp \left(\left(r - \frac{1}{2} \sigma^2 \right) (T - t) + \sigma (W_T - W_t) \right).$$

Finally, using the Brownian-increment property $W_T - W_t \sim N(0, T - t)$ (see the callout in Section 6), we may write $W_T - W_t = \sqrt{T - t} Z$ with $Z \sim N(0, 1)$, so that

$$S_T = S_t \exp \left(\left(r - \frac{1}{2} \sigma^2 \right) (T - t) + \sigma \sqrt{T - t} Z \right), \quad Z \sim N(0, 1).$$

This shows that S_T is lognormally distributed — equivalently, that $\ln S_T$ is normally distributed — with mean $\ln S_t + \left(r - \frac{1}{2} \sigma^2 \right) (T - t)$ and variance $\sigma^2 (T - t)$.

💡 Normal representation of Brownian increments

The increment of a standard Brownian motion satisfies

$$W_T - W_t \sim N(0, T - t)$$

for all $0 \leq t < T$. This is one of the defining properties of Brownian motion. If

$X \sim N(0, \sigma^2)$, then there exists a standard normal $Z \sim N(0, 1)$ such that

$$X = \sigma Z,$$

because

$$\text{Var}(\sigma Z) = \sigma^2 \text{Var}(Z) = \sigma^2.$$

Applying this fact to the Brownian increment with variance $T - t$, we write

$$W_T - W_t = \sqrt{T - t} Z, \quad Z \sim N(0, 1).$$

This is simply a reparameterisation of a normal random variable with variance $T - t$ in terms of a standard normal variable.

6. Conclusion

Combining a second-order Taylor expansion with Itô's rules yields Itô's lemma, and applying it to the value of a derivative $V(S_t, t)$ produces a diffusion whose dW_t component can be eliminated by a delta-hedged portfolio. The resulting local no-arbitrage condition is the Black-Scholes PDE Eq. (9), and evaluating the risk-neutral expectation implied by Feynman-Kac under the lognormal law of S_T delivers the closed-form call price Eq. (10). Extensions

— dividends, stochastic volatility, early exercise, jumps — follow the same three-step recipe: pick a dynamics, apply Itô's lemma, and enforce local no-arbitrage after constructing an appropriate hedged portfolio.

References

- Black, F. and Scholes, M. (1973). The Pricing of Options and Corporate Liabilities. *Journal of Political Economy*, 81(3):637–654.
- Harrison, J. M. and Pliska, S. R. (1981). Martingales and Stochastic Integrals in the Theory of Continuous Trading. *Stochastic Processes and Their Applications*, 11(3):215–260.
- Hull, J. C. (2021). *Options, Futures, and Other Derivatives*. Pearson, Boston, 11th edition.
- Itô, K. (1951). On Stochastic Differential Equations. *Memoirs of the American Mathematical Society*, 4:1–51.
- Merton, R. C. (1973). Theory of Rational Option Pricing. *The Bell Journal of Economics and Management Science*, 4(1):141–183.
- Shreve, S. E. (2004). *Stochastic Calculus for Finance II: Continuous-Time Models*. Springer, New York.